

MONTHLY MENTAL HEALTH NEWSLETTER

Presented by the Gwinnett County Sheriff's Office Mental Health Task Force



Did you know:

SEPTEMBER IS NATIONAL SUICIDE PREVENTION AWARENESS MONTH

SEPTEMBER 10TH THROUGH 16TH IS NATIONAL SUICIDE PREVENTION WEEK

SEPTEMBER 10TH IS WORLD SUICIDE PREVENTION DAY

LET'S COMBAT THIS PUBLIC HEALTH CRISIS

HELP SAVE LIVES. END THE MYTHS, AND STOP THE STIGMA WITH FACTS AND DIRECT, OPEN CONVERSATIONS.

suicidepreventionlifeline.org

MYTHS ABOUT SUICIDE (True or False)

All people who attempt suicide want to die (False)

Suicide is rare (False)

Rich people don't commit suicide (False)

Once someone attempts they will always be suicidal (False)

More women die by suicide than men (False)

People who attempt several times just want attention (False)

People who really want to commit suicide won't talk about it (False)

People who talk about suicide a lot don't really want to die and won't really try to commit suicide (False)

Asking someone if they are suicidal is dangerous because it could give them the idea to do it and then they might die (False)

When talking to someone who is suicidal we should avoid direct questions about their willingness to die. (False)

If someone who <u>was</u> suicidal suddenly becomes happy and is smilling then we know for sure they are no longer in danger of taking their own life. (False)

FACTS about SUICIDE

More people die by suicide than by homicide

More people die by suicide than by automobile accident.

While more women attempt suicide than men, more men die by suicide than women

Suicide is the second leading cause of death for people ages 10-34.

Someone dies by suicide every 11 minutes.

22 of our Veterans die by suicide daily.

Individual risk factors include:

- Previous suicide attempt(s)
- Mental illness, such as Depression
- Social isolation
- Criminal / legal problems
- Financial problems
- Impulsive or aggressive tendencies
- Job problems or Job loss
- Substance use disorder / recent withdrawal or detox from Opiates or

Other risk factors and warning signs may include:

Increased alcohol and drug use

Aggressive behavior

Withdrawal from friends, family and community

Dramatic mood swings

A family history of suicide

A history of trauma or abuse

A serious or chronic medical illness

A recent tragedy or loss / Prolonged stress

REFERENCES

The G.C.S.O. Mental Health Task Force encourages you to utilize sources such as the Substance Abuse and Mental Health Services Administration (SAMHSA) to learn more about this and other important mental health topics. www.samhsa.gov

The following sources were used in the creation of this Mental Health Newsletter;

www.nimh.nih.gov/suicideprevention

https://www.nami.org/About-Mental-Illness/ Common-with-Mental-Illness/Risk-of-Suicide



Action Steps for Helping Someone in Emotional Pain













www.nimh.nih.gov/suicideprevention

GCSO Mental Health Tasks Force

Major T. Maldonado, MA, LPC, Unit Commander 770-619-6798

Dr. D. E. Tatum, Clinical Director 770-822-3111